

# THE MOOD DISORDER QUESTIONNAIRE

Answer each of the following questions to the best of your ability, then talk with your healthcare provider.

Please see additional Important Safety Information, including **Boxed Warnings**, below and [Medication Guide](#).

 **Latuda**<sup>®</sup>  
(lurasidone HCl) tablets  
20mg | 40mg | 60mg | 80mg | 120mg

	Yes	No	
1. Has there ever been a period of time when you were not your usual self and...			
... you felt so good or so hyper that other people thought you were not your normal self or you were so hyper that you got into trouble?	<input type="checkbox"/>	<input type="checkbox"/>	
... you were so irritable that you shouted at people or started fights or arguments?	<input type="checkbox"/>	<input type="checkbox"/>	
... you felt much more self-confident than usual?	<input type="checkbox"/>	<input type="checkbox"/>	
... you got much less sleep than usual and found you didn't really miss it?	<input type="checkbox"/>	<input type="checkbox"/>	
... you were much more talkative or spoke much faster than usual?	<input type="checkbox"/>	<input type="checkbox"/>	
... thoughts raced through your head or you couldn't slow your mind down?	<input type="checkbox"/>	<input type="checkbox"/>	
... you were so easily distracted by things around you that you had trouble concentrating or staying on track?	<input type="checkbox"/>	<input type="checkbox"/>	
... you had much more energy than usual?	<input type="checkbox"/>	<input type="checkbox"/>	
... you were much more active or did many more things than usual?	<input type="checkbox"/>	<input type="checkbox"/>	
... you were much more social or outgoing than usual, for example, you telephoned friends in the middle of the night?	<input type="checkbox"/>	<input type="checkbox"/>	
... you were much more interested in sex than usual?	<input type="checkbox"/>	<input type="checkbox"/>	
... you did things that were unusual for you or that other people might have thought were excessive, foolish, or risky?	<input type="checkbox"/>	<input type="checkbox"/>	
... spending money got you or your family into trouble?	<input type="checkbox"/>	<input type="checkbox"/>	
2. If you checked YES to more than one of the above, have several of these ever happened during the same period of time?	<input type="checkbox"/>	<input type="checkbox"/>	
3. How much of a problem did any of these cause you—like being unable to work; having family, money, or legal troubles; getting into arguments or fights?			
<input type="checkbox"/> No problem	<input type="checkbox"/> Minor problem	<input type="checkbox"/> Moderate problem	<input type="checkbox"/> Serious problem

**This instrument is designed for screening purposes only and is not to be used as a diagnostic tool. Always consult with your healthcare provider.**

## IMPORTANT SAFETY INFORMATION ABOUT LATUDA

### INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS; and SUICIDAL THOUGHTS AND BEHAVIORS

Elderly people with dementia-related psychosis (having lost touch with reality due to confusion and memory loss) treated with this type of medicine are at an increased risk of death compared to patients receiving placebo (sugar pill). LATUDA is not approved for the treatment of patients with dementia-related psychosis.

Antidepressant medicines may increase suicidal thoughts or behaviors in some children, teenagers, and young adults within the first few months of treatment. Depression and other serious mental illnesses are themselves associated with an increase in the risk of suicide. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed. Report any change in these symptoms immediately to the doctor. LATUDA is not approved for use in pediatric patients with depression.

Please see additional Important Safety Information, including **Boxed Warnings**, on the next page and [Medication Guide](#).

# IMPORTANT SAFETY INFORMATION AND INDICATIONS FOR LATUDA

## **INCREASED MORTALITY IN ELDERLY PATIENTS WITH DEMENTIA-RELATED PSYCHOSIS; and SUICIDAL THOUGHTS AND BEHAVIORS**

**Elderly people with dementia-related psychosis (having lost touch with reality due to confusion and memory loss) treated with this type of medicine are at an increased risk of death compared to patients receiving placebo (sugar pill). LATUDA is not approved for the treatment of patients with dementia-related psychosis.**

**Antidepressant medicines may increase suicidal thoughts or behaviors in some children, teenagers, and young adults within the first few months of treatment. Depression and other serious mental illnesses are themselves associated with an increase in the risk of suicide. Patients on antidepressants and their families or caregivers should watch for new or worsening depression symptoms, especially sudden changes in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed. Report any change in these symptoms immediately to the doctor. LATUDA is not approved for use in pediatric patients with depression.**

LATUDA can cause serious side effects, including stroke that can lead to death, which can happen in elderly people with dementia who take medicines like LATUDA.

Neuroleptic malignant syndrome (NMS) is a rare but very serious condition that can happen in people who take antipsychotic medicines, including LATUDA. NMS can cause death and must be treated in a hospital. Call your health care provider right away if you become severely ill and have some or all of these symptoms: high fever, excessive sweating, rigid muscles, confusion, or changes in your breathing, heartbeat or blood pressure.

Tardive dyskinesia (TD) is a serious and sometimes permanent side effect reported with LATUDA and similar medicines. Tell your doctor about any movements you cannot control in your face, tongue, or other body parts, as they may be signs of TD. TD may not go away, even if you stop taking LATUDA. TD may also start after you stop taking LATUDA.

Increases in blood sugar can happen in some people who take LATUDA. Extremely high blood sugar can lead to coma or death. If you have diabetes or risk factors for diabetes (such as being overweight or a family history of diabetes), your health care provider should check your blood sugar before you start LATUDA and during therapy. Call your health care provider if you have any of these symptoms of high blood sugar (hyperglycemia) while taking LATUDA: feel very thirsty, need to urinate more than usual, feel very hungry, feel weak or tired, feel sick to your stomach, feel confused, or your breath smells fruity.

Increases in triglycerides and LDL (bad) cholesterol and decreases in HDL (good) cholesterol have been reported with LATUDA. You may not have any symptoms, so your health care provider may decide to check your cholesterol and triglycerides during your treatment with LATUDA.

Some patients may gain weight while taking LATUDA. Your doctor should check your weight regularly.

Tell your doctor if you experience any of these:

- feeling dizzy or light-headed upon standing
- decreases in white blood cells (which can be fatal)
- trouble swallowing

LATUDA and medicines like it may raise the level of prolactin. Tell your health care provider if you experience a lack of menstrual periods, leaking or enlarged breasts, or impotence.

Tell your health care provider if you have a seizure disorder, have had seizures in the past, or have conditions that increase your risk for seizures.

Tell your health care provider if you experience prolonged, abnormal muscle spasms or contractions, which may be a sign of a condition called dystonia.

LATUDA can affect your judgment, thinking, and motor skills. You should not drive or operate hazardous machinery until you know how LATUDA affects you.

LATUDA may make you more sensitive to heat. You may have trouble cooling off. Be careful when exercising or when doing things likely to cause dehydration or make you warm.

Avoid eating grapefruit or drinking grapefruit juice while you take LATUDA since these can affect the amount of LATUDA in the blood.

Tell your health care provider about all prescription and over-the-counter medicines you are taking or plan to take, since there are some risks for drug interactions with LATUDA. Tell your health care provider if you are allergic to any of the ingredients of LATUDA or take certain medications called CYP3A4 inhibitors or inducers. Ask your health care provider if you are not sure if you are taking any of these medications.

Avoid drinking alcohol while taking LATUDA.

Tell your health care provider if you are pregnant or if you are planning to get pregnant. Avoid breastfeeding while taking LATUDA.

The most common side effects of LATUDA include sleepiness or drowsiness; restlessness or feeling like you need to move around (akathisia); difficulty moving, slow movements, muscle stiffness, or tremor; runny nose/nasal inflammation, and nausea.

These are not all the possible side effects of LATUDA. For more information, ask your health care provider or pharmacist.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

## **INDICATIONS**

LATUDA is used to treat patients with:

- Depressive episodes in bipolar I disorder (bipolar depression) when used alone or with lithium or valproate in adults
- Schizophrenia in adults and adolescents 13 to 17 years of age